



Female Post Insertion Instructions

- Your insertion site has been covered with two layers of bandages. Remove the outer pressure bandage any time after 5 to 7 days. It **must** be removed as soon as it gets wet. You may replace it with a bandage to catch any anesthetic that may ooze out. The inner layer is either waterproof foam tape or steri-strips. It should be left until it falls off. If the tape or steri-strip comes off you may replace it with a band-aid.
- Do not take tub baths or get into a hot tub or swimming pool for **3 days**. You may shower but do not scrub the site until the incision is well healed (about 7 days).
- No major exercises for the incision area for the next **4 days**, this includes running, riding a horse, etc.
- The sodium bicarbonate in the anesthetic may cause the site to swell for 1-3 days. This is normal.
- The insertion site may be uncomfortable for up to 2 to 3 weeks. If there is itching or redness you may take Benadryl for relief, 50 mg. orally every 6 hours. Caution this can cause drowsiness!
- You may experience bruising, swelling, and/or redness of the insertion site which may last from a few days up to 2 to 3 weeks.
- You may notice some pinkish or bloody discoloration of the outer bandage. This is normal.
- If you experience bleeding from the incision, apply firm pressure for 5 minutes.
- Please call if you have any bleeding (not oozing) coming out of the insertion site that is not relieved by pressure.

Please call if you have any pus coming out of the insertion site, or any other signs of infection.